

**Admission LOCUS Dimension Assessment**

<b>DIMENSION</b>	<b>Level I</b>	<b>Level 2</b>	<b>Level 3</b>	<b>Level 4</b>	<b>Level 5</b>
I. Dangerousness	-Minimal Potential	-Fleeting SI or HI in past -Subst use without harm -Past self-neglect	-Sig. SI or HI without intent or plan -Past Hx of SI/HI or threats with no sig change -Past binge use with aggression towards self or others -Current self-neglect	-SI/HI & Intent or Past Hx of Attempts, without means; Able to Contract for Safety -Recent excessive subst abuse with Sig. Aggression towards self or others -Compromise of ability to care for self	-Current SI/HI & Plan & Means & clear intent or Hx of serious attempts or Command Halluc or Delusions -Repeated episodes of serious violence towards self or others while under influ of intox subst -Extreme compromise of ability to care for self or monitor environment & physical deterioration
II. Functioning	-Transient impairment secondary to exposure to clear stressor	-Freq. conflict in meaningful relationships -Mild decreased self care or neglect due to stress -Probs with ongoing social roles	-Serious conflict in Sig. Relationship -Freq. probs with ADLs with change in life rhythms -Sig decrease in role performance	-Consistent conflict in sig. Relationships with phys. Altercat -Sig. Withdrawal or avoidance -ADL compromise -Sig decr in biorhyth. -Inability to meet role expectations	-Extreme disruption in Sig Relationship(s) & Assaultive Behav with provocation -Complete withdrawal -ADL Failure -Total neglect of role responsibilities
III. Co-Morbidity	No Comorbidity	-Minor Med. Probs -Minor Subst abuse -Psych Sxs due to stress, med illness or subst abuse with no effect on primary disorder	-Med condns require med monitoring -Med condn adversely affected by 1° D/O -Subst abuse has sig impact on 1° disorder and function -Sig Psych Sxs have (-) effect on 1° D/O	-Med condns req freq med monitoring -Med cond (-) affected by primary disorder -Subst abuse threatens health & recovery -Psych Sxs impair recovery from 1° disorder.	-Sig med condn poorly controlled or life-threatening -1° disorder exac Med Condn/Uncontrolled Med Condn exac 1° disorder. -Severe uncontrolled subst abuse with w/drawal prevents recovery and or worsens 1° disorder -Acute psych sxs prevent tx or worsen 1° disorder

<b>DIMENSION</b>	<b>Level 1</b>	<b>Level 2</b>	<b>Level 3</b>	<b>Level 4</b>	<b>Level 5</b>
IVa. Stress	-No interpersonal difficulties -No major losses -Able to meet material needs; no threats to health or safety -All demands of social role easily met	-Some interper difficulties -Transition requires adjustment -Minor loss -Transient illness or injury -Conflict near home -Potential for drug or ETOH exposure -Concern over income -Performance pressure	-Interp. Discord in sig relationships -Disruption in job; legal probs -Sig loss -Decr in health status or danger -Easy access to ETOH or drugs -Increase in responsib & inability to meet obligs	-Serious disruption in family or social milieu -Pressure to join others in drug taking -Imminent incarceration, homelessness or exile -Inability to provide for physical well-being -Severe or chronic illness	-Ongoing abuse from Sig Other -Victim of, or witness to, extr violence or disaster, persecution -Sudden or unexpected death of loved one -Active exposure to & involv. in drug use -Incarceration or homelessness -Severe injury or disability -Inabil to meet basic phys needs -Chaotic & const threatening env
IVb. Support	-Multiple sources of adequate support -Effective PACT Team	-Many sources of adequate support -Support syst able to respond if called upon	-Sources of support ambivalent or limited -Resources difficult to access -Resources unable to participate fully in treatment or make adjustments -Resources only partially utilized	-Sources of support limited & unmotivated to help -Resource dysfunction or hostile to client -Existing sppts unable to meet client needs -Supports inaccessible or discouraging of assistance -Client alienated and unwilling to use sppts	-No support available
V. Recovery	-No experience with treatment and recovery -All previous treatment efforts successful	-Prev Tx successful in controlling sxs -Hx of stable recovery/ in structured setting or ongoing care	-Incomplete remission without intensive mgmt -Recovery only with prof mgmt or structured settings	-Limited response/incomplete remission, even with intensive Rx - Limited success for limited time periods	-Minimal response of Sxs and fct in highly structured settings

DIMENSION	Level 1	Level 2	Level 3	Level 4	Level 5
VI. Engagement	<ul style="list-style-type: none"> <li>-Good understanding of illness &amp; effect on function</li> <li>-Desire to make nec behavioral changes</li> <li>-(+) ther. Alliance</li> <li>-Understands rec. process &amp; personal resp</li> </ul>	<ul style="list-style-type: none"> <li>-Accepts exist. of illness &amp; willing to learn about its effects on fct</li> <li>-Willingness to change behav &amp; control sxs</li> <li>-(+) engagement in tx</li> </ul>	<ul style="list-style-type: none"> <li>-Equivocation in accept of illness</li> <li>-Limited commitment to behav change</li> <li>-Few trusting relationships</li> <li>-Won't use avail resources</li> <li>-Won't accept resp for recovery</li> </ul>	<ul style="list-style-type: none"> <li>-Rarely accepts reality of illness or disability</li> <li>-No desire to change behaviors</li> <li>-Relates poorly to tx and doesn't trust</li> <li>-Avoids contact with tx</li> <li>-Doesn't accept resp for recovery</li> </ul>	<ul style="list-style-type: none"> <li>-No awareness of illness or disability</li> <li>-Unable to understand recovery process</li> <li>-Unable to engage in tx</li> <li>-Avoidant, frightened, or guarded</li> </ul>

<b>Total Score</b>					
--------------------	--	--	--	--	--

Level of Care Utilization System for Psychiatric and Addiction Services, Adult Version 2000 American Association of Community Psychiatrists (adapted for PrimariLink use, T. W. Mandell, M.D. 2002)